

NON-HODGKIN LYMPHOMA  
Incidence

Table 1. Non-Hodgkin Lymphoma. Annual Case Counts, Age-Adjusted (2000 US Population Standard) Incidence Rates per 100,000 Population, and 95% Confidence Intervals (CI), by Sex and Race/Ethnicity, Greater Bay Area, 1988-2005

Year	TOTAL			MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI	Count	Rate	95% CI
ALL RACES									
1988	970	19.1	17.9-20.3	582	25.1	22.9-27.4	388	14.1	12.7-15.6
1989	1,023	19.6	18.4-20.9	636	26.5	24.4-28.9	387	13.6	12.3-15.1
1990	1,110	21.2	20.0-22.6	682	27.8	25.7-30.1	428	15.2	13.8-16.8
1991	1,137	21.2	20.0-22.5	721	28.7	26.6-31.1	416	14.4	13.1-15.9
1992	1,131	20.9	19.7-22.2	688	27.9	25.7-30.2	443	15.0	13.6-16.5
1993	1,105	20.1	18.9-21.3	691	27.1	25.0-29.3	414	13.9	12.6-15.3
1994	1,235	22.3	21.1-23.6	748	29.5	27.3-31.8	487	16.3	14.9-17.8
1995	1,318	23.3	22.1-24.6	832	31.4	29.3-33.7	486	15.9	14.6-17.4
1996	1,237	21.7	20.5-23.0	758	28.8	26.7-31.0	479	15.5	14.1-17.0
1997	1,192	20.5	19.3-21.7	653	24.6	22.7-26.7	539	16.9	15.5-18.4
1998	1,159	19.7	18.6-20.9	635	23.8	21.9-25.8	524	16.2	14.9-17.7
1999	1,205	20.0	18.9-21.2	671	24.8	22.9-26.8	534	16.2	14.8-17.6
2000	1,265	20.7	19.6-21.9	721	26.0	24.1-28.0	544	16.2	14.9-17.6
2001	1,348	21.7	20.6-22.9	753	26.9	25.0-29.0	595	17.6	16.2-19.1
2002	1,253	19.9	18.8-21.0	675	23.7	21.9-25.6	578	16.9	15.5-18.3
2003	1,259	19.8	18.7-20.9	715	24.9	23.1-26.9	544	15.6	14.3-16.9
2004	1,339	20.8	19.7-21.9	752	25.9	24.0-27.8	587	16.7	15.3-18.1
2005	1,269	19.5	18.5-20.7	719	24.3	22.5-26.2	550	15.5	14.2-16.9
WHITE (non-Hispanic)									
1988	746	20.7	19.3-22.3	457	28.0	25.4-30.8	289	14.4	12.8-16.2
1989	784	21.8	20.2-23.4	495	30.0	27.3-32.9	289	14.3	12.7-16.2
1990	811	22.8	21.3-24.5	492	29.9	27.3-32.8	319	16.1	14.4-18.1
1991	860	23.9	22.3-25.6	549	33.1	30.3-36.2	311	15.5	13.8-17.4
1992	844	23.1	21.6-24.8	511	30.8	28.1-33.7	333	16.2	14.5-18.1
1993	783	21.6	20.1-23.2	486	29.5	26.9-32.4	297	14.6	13.0-16.5
1994	895	24.7	23.1-26.4	546	33.2	30.4-36.2	349	17.4	15.6-19.4
1995	935	25.8	24.1-27.5	602	35.7	32.9-38.7	333	16.5	14.7-18.5
1996	851	23.5	21.9-25.2	520	31.3	28.6-34.1	331	16.5	14.7-18.4
1997	834	22.6	21.1-24.3	458	27.5	25.1-30.3	376	18.3	16.5-20.3
1998	827	22.3	20.8-23.9	476	28.4	25.9-31.1	351	16.7	15.0-18.7
1999	835	22.2	20.7-23.8	477	28.1	25.6-30.8	358	17.3	15.5-19.3
2000	864	22.9	21.4-24.5	484	28.4	25.9-31.1	380	18.2	16.4-20.2
2001	899	23.7	22.2-25.4	501	29.3	26.8-32.1	398	19.2	17.4-21.3
2002	807	21.5	20.0-23.0	444	25.6	23.3-28.2	363	17.8	16.0-19.9
2003	817	21.6	20.1-23.1	473	27.8	25.3-30.5	344	16.4	14.7-18.3
2004	865	22.9	21.4-24.6	489	28.8	26.3-31.6	376	18.1	16.2-20.1
2005	811	21.7	20.2-23.3	448	26.5	24.1-29.2	363	17.4	15.6-19.4

(Table Continues)

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Incidence

Table 1. Non-Hodgkin Lymphoma. Annual Case Counts, Age-Adjusted (2000 US Population Standard)  
(cont) Incidence Rates per 100,000 Population, and 95% Confidence Intervals (CI), by Sex and Race/Ethnicity,  
Greater Bay Area, 1988-2005

Year	TOTAL			MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI	Count	Rate	95% CI
BLACK (non-Hispanic)									
1988	44	11.8	8.4-16.5	26	15.3	9.4-25.6	18	9.2	5.4-15.1
1989	53	15.3	11.3-20.7	31	18.9	12.1-29.9	22	12.0	7.4-18.6
1990	58	15.6	11.7-20.6	38	20.2	14.1-29.7	20	10.4	6.3-16.4
1991	62	14.7	11.1-19.3	41	20.7	14.2-30.8	21	9.9	6.1-15.5
1992	63	16.1	12.2-21.1	35	19.5	13.0-29.4	28	13.2	8.7-19.5
1993	72	18.9	14.6-24.1	52	29.0	21.0-40.2	20	10.0	6.0-15.7
1994	75	17.6	13.8-22.5	59	30.1	22.5-40.8	16	7.0	4.0-11.7
1995	79	18.7	14.7-23.6	52	26.7	19.6-36.8	27	11.4	7.5-16.9
1996	73	17.7	13.8-22.5	43	21.2	15.2-30.0	30	13.9	9.3-19.9
1997	74	18.0	14.1-22.8	52	28.1	20.6-38.5	22	10.2	6.4-15.5
1998	55	13.4	10.0-17.6	30	16.3	10.7-24.7	25	11.4	7.4-16.9
1999	77	18.6	14.6-23.4	41	21.1	15.0-29.9	36	16.0	11.2-22.3
2000	62	14.8	11.3-19.1	35	18.9	13.0-27.5	27	11.5	7.6-16.8
2001	75	18.1	14.2-22.8	45	23.7	17.1-32.7	30	13.1	8.9-18.9
2002	72	17.3	13.5-21.9	43	22.2	15.9-31.0	29	12.9	8.6-18.6
2003	77	17.7	13.9-22.2	52	25.3	18.7-34.2	25	10.8	7.0-16.1
2004	71	16.6	13.0-21.1	36	17.7	12.3-25.3	35	15.2	10.5-21.3
2005	77	18.0	14.2-22.7	42	20.8	14.8-28.8	35	15.3	10.7-21.5
HISPANIC									
1988	108	22.3	18.0-27.6	62	29.5	21.4-40.5	46	17.6	12.7-23.9
1989	95	16.6	13.2-20.9	58	18.3	13.2-25.9	37	14.2	9.9-19.9
1990	129	22.5	18.5-27.4	80	26.9	20.3-35.7	49	18.2	13.4-24.5
1991	107	17.9	14.3-22.2	67	22.3	16.2-30.8	40	14.0	9.8-19.4
1992	118	19.9	16.1-24.3	78	29.5	22.4-38.9	40	12.4	8.7-17.3
1993	123	18.2	14.7-22.3	77	21.8	16.2-29.4	46	14.6	10.5-19.9
1994	121	17.7	14.4-21.7	71	18.8	13.9-25.8	50	15.9	11.6-21.3
1995	141	20.3	16.7-24.5	88	25.5	19.5-33.4	53	15.5	11.4-20.8
1996	143	21.3	17.6-25.7	91	27.9	21.5-36.2	52	15.6	11.4-20.8
1997	116	17.5	14.1-21.5	60	16.0	11.5-22.4	56	17.4	13.0-22.9
1998	121	18.7	15.2-22.9	57	18.9	13.6-26.4	64	18.5	14.0-24.1
1999	117	16.8	13.7-20.6	57	18.3	13.2-25.3	60	16.0	12.0-20.9
2000	159	20.9	17.5-24.8	89	23.9	18.4-31.1	70	18.1	13.9-23.2
2001	163	21.9	18.4-26.0	86	22.8	17.5-29.7	77	20.6	16.1-25.9
2002	168	22.3	18.9-26.4	91	26.9	21.0-34.4	77	19.0	14.9-24.0
2003	166	20.7	17.4-24.4	97	25.6	20.1-32.4	69	16.4	12.7-21.1
2004	178	21.1	17.9-24.7	99	26.2	20.7-32.9	79	17.4	13.7-21.9
2005	153	17.5	14.7-20.7	89	21.7	16.9-27.7	64	14.2	10.9-18.3

(Table Continues)

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Incidence

Table 1. Non-Hodgkin Lymphoma. Annual Case Counts, Age-Adjusted (2000 US Population Standard) Incidence Rates per 100,000 Population, and 95% Confidence Intervals (CI), by Sex and Race/Ethnicity, Greater Bay Area, 1988-2005

Year	TOTAL			MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI	Count	Rate	95% CI
ASIAN/PACIFIC ISLANDER (non-Hispanic)									
1988	61	11.7	8.7-15.7	31	12.4	8.1-19.3	30	10.8	7.0-16.3
1989	85	14.6	11.4-18.7	47	19.6	13.8-27.9	38	10.9	7.5-15.7
1990	106	15.7	12.7-19.5	67	22.2	16.8-29.5	39	10.4	7.3-14.8
1991	98	13.2	10.6-16.5	58	16.2	12.1-21.9	40	10.5	7.4-14.9
1992	98	13.8	11.1-17.3	59	19.5	14.4-26.2	39	9.5	6.6-13.5
1993	104	12.9	10.4-15.9	62	16.5	12.5-22.0	42	10.0	7.1-14.0
1994	127	16.1	13.3-19.4	63	18.7	14.1-24.7	64	14.2	10.8-18.5
1995	128	14.7	12.2-17.8	70	17.4	13.4-22.6	58	12.5	9.4-16.5
1996	133	14.0	11.7-16.8	87	20.9	16.5-26.4	46	8.7	6.3-11.8
1997	140	14.4	12.0-17.2	70	16.4	12.5-21.3	70	13.0	10.0-16.7
1998	120	11.8	9.7-14.2	54	11.3	8.4-15.2	66	12.2	9.4-15.7
1999	149	13.6	11.4-16.1	79	16.3	12.7-20.8	70	11.5	8.9-14.8
2000	161	14.0	11.9-16.5	102	19.4	15.6-23.9	59	9.6	7.2-12.5
2001	191	15.9	13.6-18.4	106	19.7	16.0-24.2	85	12.8	10.2-16.0
2002	176	14.1	12.1-16.5	78	14.3	11.2-18.2	98	14.1	11.5-17.3
2003	172	13.3	11.4-15.6	83	14.6	11.6-18.4	89	12.4	9.9-15.3
2004	197	14.4	12.5-16.7	112	18.3	15.0-22.3	85	11.3	9.0-14.1
2005	201	14.3	12.3-16.4	123	20.0	16.5-24.1	78	9.9	7.9-12.5

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Incidence

Table 2. Non-Hodgkin Lymphoma. Five-Year Case Counts, Average Annual Age-Specific Incidence Rates per 100,000 Population and 95% Confidence Intervals (CI) for All Races Combined, by Sex, 2001-2005, Greater Bay Area

Age at Diagnosis	MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI
00 - 04	7	-	-	<5	-	-
05 - 09	8	-	-	8	-	-
10 - 14	13	-	-	12	-	-
15 - 19	24	2.3	1.5-3.4	14	-	-
20 - 24	34	3.2	2.2-4.4	16	1.7	1.0-2.7
25 - 29	59	4.9	3.7-6.3	30	2.7	1.8-3.9
30 - 34	103	7.1	5.8-8.7	51	3.9	2.9-5.1
35 - 39	157	10.9	9.2-12.7	68	5.1	4.0-6.5
40 - 44	229	16.2	14.1-18.4	101	7.5	6.1-9.1
45 - 49	288	22.5	20.0-25.3	152	12.1	10.2-14.1
50 - 54	299	26.6	23.7-29.8	231	20.1	17.6-22.8
55 - 59	351	39.4	35.4-43.8	276	29.5	26.1-33.2
60 - 64	347	55.2	49.5-61.3	252	37.2	32.8-42.1
65 - 69	334	70.9	63.5-79.0	284	52.8	46.8-59.3
70 - 74	397	102.8	92.9-113.4	322	68.0	60.8-75.8
75 - 79	387	123.4	111.4-136.4	394	89.8	81.1-99.1
80 - 84	323	148.3	132.5-165.4	341	100.5	90.1-111.7
85 +	254	161.1	141.9-182.2	302	90.0	80.1-100.7
Total/Crude	3,614	22.0	21.3-22.7	2,854	17.4	16.8-18.1

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Incidence

Table 3. Non-Hodgkin Lymphoma. Five-Year Case Counts, Average Annual Age-Adjusted (2000 US Population Standard) Incidence Rates per 100,000 Population, and 95% Confidence Intervals (CI), by County, Sex, and Race/Ethnicity, Greater Bay Area, 2001-2005

County	MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI
ALL RACES						
SF Bay Area:	3,278	25.2	24.3-26.1	2,575	16.4	15.8-17.1
Alameda	739	24.7	22.9-26.6	598	16.1	14.8-17.4
Contra Costa	511	23.5	21.5-25.7	435	16.1	14.6-17.7
Marin	180	27.0	23.1-31.4	125	15.1	12.5-18.3
San Francisco	615	30.6	28.2-33.2	350	15.2	13.6-17.1
San Mateo	420	25.5	23.1-28.1	352	17.0	15.2-18.9
Santa Clara	813	23.5	21.9-25.3	715	17.7	16.4-19.0
Monterey Bay Area:	336	24.3	21.7-27.1	279	16.3	14.4-18.4
Monterey	175	22.4	19.2-26.1	138	14.6	12.3-17.3
San Benito	21	18.4	11.1-29.7	29	25.1	16.7-36.5
Santa Cruz	140	28.1	23.5-33.5	112	17.2	14.1-20.8
WHITE (non-Hispanic)						
SF Bay Area:	2,105	27.7	26.5-28.9	1,643	17.8	17.0-18.8
Alameda	404	26.3	23.7-29.0	348	18.2	16.3-20.3
Contra Costa	366	23.9	21.5-26.6	303	16.0	14.2-18.0
Marin	156	26.9	22.7-31.9	114	15.2	12.5-18.7
San Francisco	376	38.7	34.8-43.4	179	17.5	14.9-20.9
San Mateo	295	28.0	24.8-31.5	248	19.2	16.8-22.1
Santa Clara	508	26.2	23.9-28.7	451	19.0	17.3-20.9
Monterey Bay Area:	250	26.4	23.2-30.0	201	17.1	14.7-19.8
Monterey	122	25.5	21.1-30.6	94	15.7	12.7-19.7
San Benito	13	-	-	17	23.6	13.6-39.3
Santa Cruz	115	28.3	23.2-34.3	90	17.3	13.9-21.7
BLACK (non-Hispanic)						
SF Bay Area:	210	22.2	19.2-25.6	152	13.6	11.5-16.0
Alameda	92	21.6	17.3-26.7	74	13.6	10.7-17.2
Contra Costa	29	17.7	11.5-26.8	33	14.7	10.0-21.0
Marin	<5	-	-	<5	-	-
San Francisco	58	37.8	28.7-49.3	30	17.2	11.4-25.3
San Mateo	12	-	-	7	-	-
Santa Clara	17	13.3	7.7-30.5	7	-	-
Monterey Bay Area:	8	-	-	<5	-	-
Monterey	8	-	-	<5	-	-
San Benito	<5	-	-	<5	-	-
Santa Cruz	<5	-	-	<5	-	-

(Table Continues)

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Incidence

Table 3. Non-Hodgkin Lymphoma. Five-Year Case Counts, Average Annual Age-Adjusted (2000 US Population Standard) Incidence Rates per 100,000 Population, and 95% Confidence Intervals (CI), by County, Sex, and Race/Ethnicity, Greater Bay Area, 2001-2005

County	MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI
HISPANIC						
SF Bay Area:	404	25.8	23.0-28.9	312	17.4	15.5-19.6
Alameda	96	29.3	23.2-36.8	66	16.1	12.3-20.7
Contra Costa	49	21.7	15.2-30.7	47	17.3	12.5-23.4
Marin	10	-	-	5	-	-
San Francisco	74	34.2	26.4-44.0	46	19.3	14.1-25.9
San Mateo	43	19.1	13.3-27.3	45	18.7	13.6-25.3
Santa Clara	132	24.6	20.0-30.2	103	18.0	14.5-22.0
Monterey Bay Area:	58	19.1	13.7-26.5	54	16.8	12.4-22.7
Monterey	36	18.1	11.7-27.9	31	14.9	9.7-22.2
San Benito	8	-	-	8	-	-
Santa Cruz	14	-	-	15	20.5	10.9-36.2
ASIAN/PACIFIC ISLANDER (non-Hispanic)						
SF Bay Area:	488	17.6	16.0-19.3	420	12.0	10.9-13.2
Alameda	121	19.3	15.8-23.4	89	11.2	9.0-13.9
Contra Costa	47	18.4	13.2-25.2	43	13.8	9.9-18.8
Marin	9	-	-	<5	-	-
San Francisco	103	15.4	12.5-18.8	92	11.0	8.8-13.8
San Mateo	62	21.0	15.8-27.7	48	11.2	8.2-15.1
Santa Clara	146	16.4	13.7-19.6	145	14.3	12.0-17.0
Monterey Bay Area:	14	-	-	15	12.2	6.8-21.1
Monterey	8	-	-	9	-	-
San Benito	<5	-	-	<5	-	-
Santa Cruz	6	-	-	<5	-	-

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Mortality

Table 4. Non-Hodgkin Lymphoma. Annual Death Counts, Age-Adjusted (2000 US Population Standard) Mortality Rates per 100,000 Population, and 95% Confidence Intervals (CI), by Sex and Race/Ethnicity, Greater Bay Area, 1988-2005

Year	TOTAL			MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI	Count	Rate	95% CI
ALL RACES									
1988	392	8.0	7.2-8.9	221	10.6	9.2-12.2	171	6.1	5.2-7.1
1989	368	7.6	6.9-8.5	198	9.9	8.5-11.5	170	6.0	5.1-7.0
1990	383	7.8	7.0-8.6	223	10.6	9.2-12.2	160	5.6	4.8-6.6
1991	385	7.7	6.9-8.5	208	9.6	8.3-11.2	177	6.1	5.2-7.1
1992	422	8.3	7.5-9.1	228	10.9	9.5-12.5	194	6.5	5.6-7.5
1993	422	8.1	7.4-9.0	228	10.5	9.1-12.1	194	6.5	5.6-7.5
1994	434	8.3	7.5-9.1	221	10.1	8.8-11.6	213	7.0	6.1-8.0
1995	465	8.7	7.9-9.5	259	11.4	10.0-12.9	206	6.7	5.8-7.7
1996	422	7.7	7.0-8.5	217	9.1	7.9-10.5	205	6.5	5.7-7.5
1997	513	9.1	8.4-10.0	278	11.2	9.9-12.6	235	7.3	6.4-8.3
1998	455	8.0	7.3-8.7	232	9.4	8.2-10.7	223	6.8	5.9-7.7
1999	415	7.2	6.5-7.9	243	10.0	8.7-11.3	172	5.2	4.4-6.0
2000	464	7.8	7.1-8.6	231	9.3	8.1-10.6	233	6.7	5.9-7.7
2001	427	7.1	6.4-7.8	235	9.3	8.1-10.6	192	5.5	4.8-6.4
2002	460	7.5	6.9-8.3	250	9.6	8.4-10.9	210	5.9	5.2-6.8
2003	447	7.3	6.6-8.0	255	9.9	8.7-11.2	192	5.4	4.6-6.2
2004	461	7.3	6.7-8.0	252	9.4	8.3-10.7	209	5.8	5.0-6.7
2005	425	6.7	6.1-7.4	232	8.7	7.6-9.9	193	5.4	4.6-6.2
WHITE (non-Hispanic)									
1988	317	8.9	7.9-9.9	174	11.6	9.9-13.6	143	6.8	5.7-8.1
1989	288	8.2	7.3-9.2	153	10.7	9.0-12.6	135	6.4	5.4-7.7
1990	314	8.9	7.9-9.9	180	12.1	10.3-14.1	134	6.4	5.3-7.6
1991	303	8.5	7.6-9.6	159	10.7	9.0-12.6	144	6.8	5.7-8.1
1992	317	8.9	7.9-9.9	167	11.5	9.8-13.5	150	7.1	6.0-8.4
1993	320	8.9	7.9-9.9	164	11.0	9.3-12.9	156	7.3	6.2-8.6
1994	336	9.3	8.3-10.3	168	11.3	9.6-13.2	168	7.9	6.7-9.2
1995	358	9.7	8.7-10.8	198	12.6	10.9-14.6	160	7.5	6.3-8.8
1996	320	8.7	7.8-9.7	162	10.3	8.7-12.0	158	7.4	6.2-8.7
1997	361	9.8	8.8-10.8	196	12.1	10.4-13.9	165	7.6	6.5-9.0
1998	349	9.2	8.3-10.3	183	11.4	9.8-13.3	166	7.4	6.3-8.7
1999	300	7.9	7.0-8.9	177	11.1	9.5-12.9	123	5.5	4.5-6.6
2000	326	8.3	7.5-9.3	157	9.8	8.3-11.5	169	7.4	6.3-8.7
2001	309	7.9	7.1-8.9	165	10.0	8.6-11.7	144	6.4	5.3-7.6
2002	310	8.0	7.1-9.0	176	10.8	9.3-12.6	134	5.8	4.8-7.0
2003	318	8.2	7.3-9.2	180	11.2	9.6-13.0	138	6.1	5.1-7.3
2004	301	7.7	6.9-8.7	165	10.1	8.6-11.8	136	6.0	5.0-7.2
2005	285	7.3	6.5-8.3	158	9.7	8.2-11.4	127	5.7	4.7-6.9

(Table Continues)

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Mortality

Table 4. Non-Hodgkin Lymphoma. Annual Death Counts, Age-Adjusted (2000 US Population Standard) (cont) Mortality Rates per 100,000 Population, and 95% Confidence Intervals (CI), by Sex and Race/Ethnicity, Greater Bay Area, 1988-2005

Year	TOTAL			MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI	Count	Rate	95% CI
BLACK (non-Hispanic)									
1988	20	5.5	3.3-9.0	14	-	-	6	-	-
1989	23	7.4	4.6-11.7	14	-	-	9	-	-
1990	15	4.3	2.4-7.6	10	-	-	5	-	-
1991	15	4.3	2.3-7.5	8	-	-	7	-	-
1992	28	8.4	5.5-12.6	16	12.1	6.6-21.5	12	-	-
1993	18	5.4	3.2-8.9	7	-	-	11	-	-
1994	23	6.3	3.9-9.7	10	-	-	13	-	-
1995	24	7.0	4.5-10.7	10	-	-	14	-	-
1996	17	4.5	2.6-7.4	9	-	-	8	-	-
1997	36	9.1	6.4-12.8	23	12.9	8.0-20.7	13	-	-
1998	23	6.0	3.8-9.1	9	-	-	14	-	-
1999	18	4.8	2.8-7.7	8	-	-	10	-	-
2000	26	6.5	4.2-9.7	12	-	-	14	-	-
2001	22	5.6	3.5-8.6	13	-	-	9	-	-
2002	24	6.3	4.0-9.5	13	-	-	11	-	-
2003	15	3.8	2.1-6.4	11	-	-	<5	-	-
2004	34	8.5	5.9-12.1	19	10.1	6.0-16.5	15	6.9	3.8-11.5
2005	18	4.6	2.7-7.5	8	-	-	10	-	-
HISPANIC									
1988	25	5.7	3.6-9.0	13	-	-	12	-	-
1989	32	7.0	4.7-10.4	18	9.2	5.0-16.3	14	-	-
1990	26	5.6	3.5-8.6	14	-	-	12	-	-
1991	33	7.2	4.8-10.5	17	7.6	4.2-13.5	16	6.4	3.6-10.7
1992	31	5.7	3.8-8.4	16	7.6	4.0-13.8	15	4.6	2.6-8.0
1993	47	9.2	6.6-12.6	33	16.2	10.6-24.3	14	-	-
1994	32	6.3	4.2-9.3	18	8.2	4.4-14.5	14	-	-
1995	38	7.0	4.8-10.0	21	10.0	5.6-17.0	17	5.5	3.1-9.1
1996	44	8.1	5.8-11.2	23	9.5	5.7-15.6	21	6.9	4.2-10.8
1997	56	9.9	7.3-13.2	27	9.8	6.0-15.8	29	9.5	6.3-13.9
1998	34	6.2	4.2-9.0	14	-	-	20	6.4	3.8-10.1
1999	51	9.3	6.8-12.4	31	14.0	9.0-21.1	20	6.3	3.8-10.0
2000	41	6.8	4.8-9.5	20	7.6	4.3-12.9	21	6.3	3.9-9.8
2001	43	6.9	4.9-9.5	25	8.7	5.3-13.8	18	5.4	3.2-8.7
2002	56	8.6	6.4-11.4	30	9.8	6.4-15.0	26	7.4	4.8-11.0
2003	49	6.7	4.8-9.1	31	9.3	6.0-14.2	18	4.6	2.7-7.4
2004	57	7.7	5.7-10.1	30	8.5	5.5-12.9	27	6.8	4.5-10.0
2005	46	6.3	4.6-8.5	27	8.5	5.5-13.0	19	4.7	2.8-7.4

(Table Continues)

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Mortality

Table 4. Non-Hodgkin Lymphoma. Annual Death Counts, Age-Adjusted (2000 US Population Standard) Mortality Rates per 100,000 Population, and 95% Confidence Intervals (CI), by Sex and Race/Ethnicity, Greater Bay Area, 1988-2005

Year	TOTAL			MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI	Count	Rate	95% CI
ASIAN/PACIFIC ISLANDER (non-Hispanic)									
1988	27	5.2	3.3-8.1	17	6.7	3.8-12.3	10	-	-
1989	25	4.8	2.9-7.6	13	-	-	12	-	-
1990	26	4.6	2.9-7.1	17	6.6	3.7-11.7	9	-	-
1991	34	5.0	3.4-7.4	24	8.1	5.1-13.0	10	-	-
1992	44	6.6	4.7-9.2	28	9.0	5.8-13.8	16	4.5	2.5-7.8
1993	36	5.3	3.6-7.7	24	8.2	5.0-12.9	12	-	-
1994	43	5.5	3.9-7.7	25	7.6	4.8-11.9	18	3.9	2.3-6.6
1995	45	5.9	4.2-8.1	30	8.8	5.8-13.1	15	3.7	2.0-6.4
1996	40	4.6	3.3-6.5	22	5.9	3.6-9.5	18	3.7	2.2-6.2
1997	60	6.9	5.2-9.1	32	8.2	5.4-12.0	28	5.9	3.9-8.8
1998	48	4.8	3.5-6.6	25	5.3	3.4-8.2	23	4.4	2.7-6.8
1999	46	4.3	3.1-5.9	27	6.1	4.0-9.3	19	3.0	1.8-4.9
2000	69	6.5	5.0-8.4	42	8.2	5.8-11.4	27	5.0	3.3-7.4
2001	51	4.6	3.4-6.2	30	6.6	4.4-9.7	21	3.2	2.0-5.0
2002	70	5.9	4.6-7.6	31	6.1	4.1-9.0	39	5.8	4.1-8.0
2003	64	5.5	4.2-7.0	32	6.4	4.4-9.3	32	4.8	3.3-6.8
2004	69	5.5	4.3-7.0	38	7.1	5.0-9.9	31	4.3	2.9-6.2
2005	75	5.6	4.4-7.1	38	6.8	4.8-9.5	37	4.8	3.4-6.7

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Mortality

Table 5. Non-Hodgkin Lymphoma. Five-Year Death Counts, Average Annual Age-Specific Mortality Rates per 100,000 Population and 95% Confidence Intervals (CI) for All Races Combined, by Sex, 2001-2005, Greater Bay Area

Age at Diagnosis	MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI
00 - 04	<5	-	-	<5	-	-
05 - 09	<5	-	-	<5	-	-
10 - 14	<5	-	-	<5	-	-
15 - 19	<5	-	-	<5	-	-
20 - 24	<5	-	-	<5	-	-
25 - 29	12	-	-	<5	-	-
30 - 34	5	-	-	9	-	-
35 - 39	15	1.0	0.6-1.7	8	-	-
40 - 44	32	2.3	1.5-3.2	15	1.1	0.6-1.8
45 - 49	43	3.4	2.4-4.5	21	1.7	1.0-2.6
50 - 54	70	6.2	4.9-7.9	39	3.4	2.4-4.6
55 - 59	85	9.6	7.6-11.8	49	5.2	3.9-6.9
60 - 64	109	17.3	14.2-20.9	56	8.3	6.3-10.7
65 - 69	118	25.1	20.7-30.0	76	14.1	11.1-17.7
70 - 74	147	38.1	32.2-44.7	122	25.8	21.4-30.8
75 - 79	187	59.7	51.4-68.8	189	43.1	37.1-49.7
80 - 84	213	97.8	85.1-111.8	183	53.9	46.4-62.3
85 +	180	114.2	98.1-132.1	221	65.9	57.5-75.1
Total/Crude	1,224	7.4	7.0-7.9	996	6.1	5.7-6.5

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Mortality

Table 6. Non-Hodgkin Lymphoma. Five-Year Death Counts, Average Annual Age-Adjusted (2000 US Population Standard) Mortality Rates per 100,000 Population, and 95% Confidence Intervals (CI), by County, Sex, and Race/Ethnicity, Greater Bay Area, 2001-2005

County	MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI
ALL RACES						
SF Bay Area:	1,104	9.4	8.8-10.0	889	5.5	5.2-5.9
Alameda	245	9.1	8.0-10.3	185	5.0	4.3-5.8
Contra Costa	180	9.2	7.9-10.7	178	6.4	5.5-7.4
Marin	81	12.8	10.2-16.1	38	4.7	3.3-6.7
San Francisco	176	9.6	8.2-11.1	127	5.0	4.1-6.0
San Mateo	174	11.0	9.4-12.8	125	5.8	4.8-7.0
Santa Clara	248	8.1	7.1-9.2	236	5.9	5.2-6.7
Monterey Bay Area:	120	9.3	7.6-11.1	107	6.1	5.0-7.4
Monterey	71	9.4	7.3-11.9	47	4.9	3.6-6.5
San Benito	5	-	-	6	-	-
Santa Cruz	44	9.9	7.1-13.4	54	8.1	6.0-10.7
WHITE (non-Hispanic)						
SF Bay Area:	756	10.4	9.7-11.2	595	5.9	5.5-6.5
Alameda	142	9.8	8.3-11.6	113	5.5	4.5-6.7
Contra Costa	141	9.9	8.3-11.7	140	6.7	5.6-8.0
Marin	74	12.9	10.1-16.6	37	5.0	3.5-7.4
San Francisco	103	12.2	10.0-15.4	58	5.0	3.7-7.2
San Mateo	133	12.4	10.4-14.9	89	6.1	4.9-7.9
Santa Clara	163	8.9	7.6-10.5	158	6.3	5.3-7.4
Monterey Bay Area:	88	9.5	7.6-11.9	84	6.4	5.1-8.1
Monterey	49	10.1	7.5-13.6	37	5.4	3.8-7.9
San Benito	<5	-	-	<5	-	-
Santa Cruz	38	10.0	7.0-14.0	43	7.4	5.3-10.5
BLACK (non-Hispanic)						
SF Bay Area:	62	7.5	5.7-9.8	48	4.4	3.2-5.9
Alameda	29	7.6	5.0-11.1	23	4.1	2.6-6.3
Contra Costa	11	-	-	13	-	-
Marin	<5	-	-	<5	-	-
San Francisco	13	-	-	8	-	-
San Mateo	<5	-	-	<5	-	-
Santa Clara	5	-	-	<5	-	-
Monterey Bay Area:	<5	-	-	<5	-	-
Monterey	<5	-	-	<5	-	-
San Benito	<5	-	-	<5	-	-
Santa Cruz	<5	-	-	<5	-	-

(Table Continues)

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.

NON-HODGKIN LYMPHOMA  
Mortality

Table 6. Non-Hodgkin Lymphoma. Five-Year Death Counts, Average Annual Age-Adjusted (2000 US Population Standard) Mortality Rates per 100,000 Population, and 95% Confidence Intervals (CI), by County, Sex, and Race/Ethnicity, Greater Bay Area, 2001-2005

County	MALES*			FEMALES		
	Count	Rate	95% CI	Count	Rate	95% CI
HISPANIC						
SF Bay Area:	117	9.1	7.4-11.1	91	5.7	4.5-7.0
Alameda	30	10.8	7.0-16.1	21	5.9	3.6-9.1
Contra Costa	15	7.6	3.9-14.3	12	-	-
Marin	<5	-	-	<5	-	-
San Francisco	17	10.0	5.7-16.5	14	-	-
San Mateo	18	10.2	5.8-17.3	15	7.0	3.9-11.7
Santa Clara	35	7.7	5.2-11.4	29	5.6	3.7-8.2
Monterey Bay Area:	26	8.2	4.9-13.6	17	6.3	3.5-10.5
Monterey	17	8.0	4.1-15.5	7	-	-
San Benito	<5	-	-	<5	-	-
Santa Cruz	5	-	-	9	-	-
ASIAN/PACIFIC ISLANDER (non-Hispanic)						
SF Bay Area:	165	6.7	5.7-7.9	155	4.7	3.9-5.5
Alameda	42	7.6	5.4-10.5	28	3.7	2.4-5.4
Contra Costa	11	-	-	13	-	-
Marin	<5	-	-	<5	-	-
San Francisco	43	6.5	4.7-8.9	47	5.0	3.7-6.9
San Mateo	21	8.7	5.2-13.8	18	4.6	2.7-7.4
Santa Clara	45	6.0	4.3-8.2	48	5.3	3.9-7.1
Monterey Bay Area:	<5	-	-	5	-	-
Monterey	<5	-	-	<5	-	-
San Benito	<5	-	-	<5	-	-
Santa Cruz	<5	-	-	<5	-	-

\*Veterans Health Administration hospitals did not report cancer cases in 2005. Therefore, case counts and incidence rates for adult males in 2005 are underestimated and should be interpreted with caution.